

Global Embedding of Analytic Branes into Einstein MD Bulk Cosmology

Nikolaos I. Katzourakis*

University of Athens, Department of Mathematics

e-mail: nkatzourakis@math.uoa.gr

Abstract

In recent work [26], [27] we have shown that *any n -dimensional analytic semi-Riemannian manifold $\mathcal{M}$ has a global analytic and isometric embedding into an Einstein manifold $\mathcal{E}$ of dimension $n + d$, $d \geq 1$ ($\Lambda \in \mathbb{R}$).*

In addition, if $\mathcal{M}$ is an analytic spacetime manifold, there exists an Einstein global embedding space $\mathcal{E}^\Sigma$, admitting unsmoothable differential-topological singularities ($0 - D$ points or extended $(d - k) - D$ spatial "anomalies") living into the complement $\mathcal{E}^\Sigma \setminus \mathcal{M}$. They were expounded as wormholes.

In this paper we continue extending this result on Cosmological Brane embeddings:

Any finite-many analytic semi-Riemannian manifolds $\mathcal{M}_1^{(n_1)}, \dots, \mathcal{M}_m^{(n_m)}$ have all a global isometric analytic embedding into an $(\sum_{1 \leq k \leq m} n_k + 1)$ -Einstein Bulk $\mathcal{E}$ as disjoint submanifolds, while $\pi_m(\mathcal{E})$ splits.

Keywords: Spacetime Embedding, Einstein space, Campbell-Magaard Theorem; Brane-World, MD Gravity, Brane Cosmology

MSC Subject Classification 2000: Primary 53C20, 53C25, 32C09, 53Cxx, Secondary 83C75, 83C55, 83E15, 83F05.

*Address: Troias 6, 18541, Kaminia, Pireaus, Greece

Introduction

Multi-dimensional theories of gravity and, in particular, the recent Braneworld Cosmological scenario, in which the Universe of our perception is a submanifold (viz. *(mem)-Brane*) embedded in a multi-dimensional (MD) manifold (viz. *Bulk*), have generated a marvelous interest in mathematical and physical aspects of embedding theories based on the 4-spacetime of *General Relativity* (*G.T.R.*). The theme of the present paper is devoted to the *mathematical* perspective of this subject.

According to the G.T.R., the model we adopt to describe the large scale structure of the Universe is a 4-manifold equipped with a *semi-Riemannian metric* structure, the solution of the *field equations* which (the latter) correlate the *Curvature* of spacetime with the distribution of *matter*, the *energy* and the *momentum* ([5], [7], [12], [14]).

The possibility that there may exist more than $3 + 1$ space-time dimensions was taken into account not long after the final formulation of G.T.R.. *Kaluza* and *Klein* (K.-K.) *unified 4D vacuum Relativity* and classical *electromagnetism* ([15]) on a 5-manifold that satisfies the vacuum field equations (curved by the contribution of the photons). The extra dimension was considered topologically *compact* (*homeomorphic to S^1*), *infinitesimal* (thus invisible) and the 4-spacetime was assumed not to depend on the 5th dimension. In fact, the Universe was considered as a *hyper-cylinder*, locally *homeomorphic to $\mathbb{R}^4 \times S^1$* .

In a modern approach called *Space-Time-Matter* or *Induced Matter Theory* (S.T.M.) ([17], [18], [15] and references therein) these assumptions are removed and the result is the *unification of Relativity and electromagnetism with their sources* (*mass* and *charge*), expounded (the latter) as an artifact induced on the 4-manifold, generated by the (non-linear) geometry of invisible extra spatial dimensions.

One of the most recent Cosmological scenarios concerning the geometric structure of the (classical) Universe introduced by L. Randall and R. Sundrum ([37], [38]) claims that the Universe of our perception is a (mem-)Brane of $3 + 1$ dimensions embedded in a multi-dimensional Bulk of at least 5 dimensions. The main ansatz for this (possibly moving in the Bulk) Brane is that *matter is confined on*

the Brane and only gravity and electromagnetic fields can propagate on the bulk.

Mathematically, these theories are modeled on a local embedding theorem of Riemannian Geometry, due to Campbell who was the first who stated it [2], and Magaard ([8], [30]) who presented a strict proof (see [34], [39], [43]): *any analytic n -manifold can be locally and isometrically embedded into some other Ricci-flat of $n+1$ dimensions ($R_{AB} = 0$).* In a sense, this is equivalent to the *local embedding of Einstein 4D Relativity, into some 5D Relativity in the vacuum where no matter exists* ("wood", according to Einstein), but only *geometrical properties* ("marble").

It becomes clear that in the light of these theories that are entirely based on differential-geometric (though *local* only) embeddings, new *global results are needed*, in order to provide the *supporting mathematical framework for Brane-Worlds*. This owes to the fact that the starting point of an MD gravitational-cosmological theory is the 4-dimensional G.T.R. that has to be incorporated in the new context.

In recent work ([26], [27]) the author proved that **the Campbell-Magaard (C.-M.) Theorem** (extended by *Dahia & Romero* in [17] on *Einstein from Ricci-flat spaces*) **holds true globally**:

Any n -dimensional analytic semi-Riemannian manifold $\mathcal{M}$ has a global analytic isometric embedding into an Einstein manifold $\mathcal{E}$ of dimension $n + d$, $d \geq 1$ ($\Lambda \in \mathbb{R}$).

This $\mathcal{E}$ may admit (if desired) 1 for embedding codimension $d = 1$ & countable many for $d > 1$ unsmoothable differential-topological singularities, which may be distinct points into spacetime or extended spatial anomalies ([27]). These pathologies living into the complement $\mathcal{E} \setminus \mathcal{M}$ were expounded as *wormholes connecting distant points of spacetime*.

In this sequel we continue showing that *any finite many semi-Riemannian manifolds $\mathcal{M}_1, \dots, \mathcal{M}_m$ have all a global isometric analytic embedding into an Einstein space $\mathcal{E}$ as disjoint analytic submanifolds. The minimum dimensionality of $\mathcal{E}$ is the sum of the respective of the $\mathcal{M}_k$'s plus 1 and this is optimum.*

The process in the proof is a slight modification of that presented in [26]. We form a bulk that contains all the $\mathcal{M}_k$'s as disjoint submanifolds and using the

known local embedding results and certain differential-geometric & topological arguments we obtain the requested global extension.

In *Section 1* we give the necessary results we will use throughout the rest of the paper. *Section 2* contains the proof of the basic result stated above. In *Section 3* we study the topology of such embedding Einstein spaces, giving a Homotopic description of the Einstein manifolds $\mathcal{E}$ we construct, which proves to be quite simple, due to direct sum decomposition: $\pi_q(\mathcal{E})$ can be completely determined if $\pi_q(\mathcal{M}_1), \dots, \pi_q(\mathcal{M}_m)$ are known.

Notations: In the sequel, the terms **Spacetime**, **Brane**, **Bulk** will be used in the place of *paracompact*, *Hausdorff*, *connected manifolds (finite dimensional)*, equipped with a *semi-Riemannian metric structure* and the *corresponding Levi-Civita linear connection*.

1. Preliminaries

In this section we collect for the readers convenience some fundamental notions retrieved from the references that will be employed in the rest of the paper.

Definition 1.1. The *G.T.R. field equations* on a semi-Riemannian manifold $(\mathcal{M}^{(n)}; \nabla, g)$ in presence of *Cosmological Constant* $\Lambda \in \mathbb{R}$ are:

$$G \equiv Ric^\nabla(g) - \frac{1}{2}R^{(n)}g = kT - \Lambda g$$

where $Ric, g, T \in \Gamma(T^*\mathcal{M} \otimes T^*\mathcal{M})$, $k \in \mathbb{R}$, G is the *Einstein tensor of Curvature* and $R^{(n)}$ is the *scalar Curvature*, defined as the double contraction with the metric. T is called the *Stress-Energy-Momentum tensor*.

Definition 1.2. A semi-Riemannian manifold $(\mathcal{M}^{(n)}; \nabla^{\mathcal{M}}, g_{\mathcal{M}})$ is said to be an *Einstein space* if there exists a $\Lambda \in \mathbb{R}$ such that:

$$Ric^{\nabla^{\mathcal{M}}}(g_{\mathcal{M}}) = \frac{2\Lambda}{n-2} g_{\mathcal{M}}$$

That is, its metric is a solution of the previous equations with $T = 0$. If $\Lambda = 0$ the manifold is said to be *Ricci-flat*.

The following has been originally retrieved from reference [17] and used also in [26], [27], as the local analogues of the basic results.

Theorem 1.3. (*Campbell-Magaard-Dahia-Romero local embedding into Einstein Spaces*)

Let $(\mathcal{M}^{(n)}, g_{\mathcal{M}})$ be a semi-Riemannian analytic manifold and

$$g|_U = g_{ij} dx^i \otimes dx^j \in \Gamma_{\alpha_0}(T^*\mathcal{M} \otimes T^*\mathcal{M})$$

a local analytic at $0 \equiv \phi(p) \in \mathbb{R}^n$ presentation of its metric, with respect to $(U, \phi) \in \mathcal{A}_{\mathcal{M}}$.

Then, there exists a local isometric embedding (analytic at $\phi(p)$) of U into some (coordinated) semi-Riemannian manifold $(\mathcal{N}^{(n+1)}; g_{\mathcal{N}}, \nabla^{\mathcal{N}})$:

$$\varepsilon : U \longrightarrow \varepsilon(U) \subseteq \mathcal{N}$$

which is an Einstein space in the local embedding domain:

$$Ric^{\nabla^{\varepsilon(U)}}(g_{\mathcal{N}}|_{\varepsilon(U)}) = \frac{2\Lambda}{n-1} g_{\mathcal{N}}|_{\varepsilon(U)} \in \Gamma(T^*\varepsilon(U) \otimes T^*\varepsilon(U))$$

that is

$$R_{AB} dx^A \otimes dx^B = \frac{2\Lambda}{n-1} g_{AB} dx^A \otimes dx^B$$

Setting $\Lambda = 0$, Th. 1.3 reduces to the Campbell-Magaard Theorem of embedding into Ricci-flat spaces ([8],[26], [27] etc).

The following can be easily deduced from Th. 1.3 and the Criterion of local isometric embeddings (see [8], [17], [26], [27]):

Corollary 1.4. Let $(U, \chi = \{x^A\}_{1 \leq A \leq n+1})$ be a patch of an analytic $(n+1)$ -manifold $\mathcal{M}$ with local metric of the diagonal form ($g_{a(n+1)} \equiv 0$):

$$g_U = g_{ab} dx^a \otimes dx^b + \psi dx^{n+1} \otimes dx^{n+1}$$

Then, if the functions g_{ab} are analytic at $0 \in \chi(U) \subseteq \mathbb{R}^{n+1}$, there exists an analytic at 0 function

$$\psi \in C_{\alpha_0}^\infty(U \longrightarrow \mathbb{R})$$

such that the coordinated domain U to be a Einstein space, with respect to g_U and the respective Levi-Civita metric connection ∇^U .

2. Global multi-Brane Embedding into Einstein Manifolds

The manifolds we employ in this section are considered real analytic. Notwithstanding, it follows from results of Whitney [44] and Grauert [24] that this is not restrictive, provided that any C^k or smooth (C^∞) manifold has a C^k (resp. C^∞) diffeomorphism with an analytic one, that is, it *carries a unique compatible real analytic structure*. See also [10].

(*This Theorem 2.1 is almost a replica of the respective embedding results of [26] & [27]. Notwithstanding, for complementary reasons, we reformulate a self contained proof from the very beginning, instead of giving citations to the prequels.*)

We state now our basic result:

Theorem 2.1. (*Global Isometric Embedding of Branes into Einstein manifolds in codimension 1*)

Any n_k -dimensional analytic semi-Riemannian manifolds $(\mathcal{M}_k; \nabla^{\mathcal{M}_k}, g_{\mathcal{M}_k})$, $1 \leq k \leq m$ (Branes) have all a global isometric analytic embedding into an $(\sum_{k=1}^m n_k + 1)$ -dimensional Einstein manifold $(\mathcal{E}; \nabla^{\mathcal{E}}, g_{\mathcal{E}})$ satisfying:

$$\text{Ric}^{\nabla^{\mathcal{E}}}(g_{\mathcal{E}}) = \frac{2\Lambda}{\left(\sum_{k=1}^m n_k\right) - 1} g_{\mathcal{E}}$$

as disjoint submanifolds for any $\Lambda \in \mathbb{R}$.

Proof. Step 1: Construction of the Bulk $\mathcal{E}$ that contains disjoint copies of $\mathcal{M}_1, \dots, \mathcal{M}_m$.

Let $\mathcal{F}$ be an abstract analytic curve (1-dimensional manifold). We set

$$\mathcal{M} := \mathcal{M}_1 \times \dots \times \mathcal{M}_m$$

and $\mathcal{A}_{\mathcal{M}}$ will denote the Atlas of differential structure of this product manifold. This product contains as submanifolds the given $\mathcal{M}_k$'s (each inclusion map is an embedding) in the induced structure. For convenience, we set

$$n := \sum_{k=1}^m n_k \in \mathbb{N}$$

We define the fibre bundles over $\mathcal{M}$ & $\mathcal{F}$:

$$\pi : \mathcal{E} \longrightarrow \mathcal{M}$$

$$pr_{\mathcal{F}} : \mathcal{E} \longrightarrow \mathcal{F}$$

where

$$\mathcal{E} := \mathcal{M}_1 \times \dots \times \mathcal{M}_m \times \mathcal{F} \equiv \mathcal{M} \times \mathcal{F}$$

and

$$\pi := pr_{\mathcal{M}} : \{(p_1, \dots, p_m)\} \times \mathcal{F} \longmapsto (p_1, \dots, p_m) \equiv p \in \mathcal{M}$$

This construction gives a Bulk $\mathcal{E}$ that contains all the $\mathcal{M}_k$'s as submanifolds. We shall treat $\mathcal{M}$ as if it was a single manifold ignoring the product structure.

The local (coordinated) trivializations have the form:

$$\pi^{-1}(U)|_V \xrightarrow[id]{\cong} U \times (\mathcal{F}|_V) \equiv W \in \mathcal{A}_{\mathcal{E}}$$

where $(U; \{x^a\}_{1 \leq a \leq n}) \in \mathcal{A}_{\mathcal{M}}$, $(V; y) \in \mathcal{A}_{\mathcal{F}}$, the atlases of differential structure. In addition, the $\mathcal{M}_k$'s are all disjoint submanifolds of $\mathcal{E}$: to see this, it suffices to chose m -many distinct points $f_1, \dots, f_m$ of the fibre $\mathcal{F}$ and identify each of the $\mathcal{M}_k$'s with a copy living into the bulk $\mathcal{E}$

$$\mathcal{M}_k \equiv \underbrace{\emptyset \times \dots \times \emptyset \times \mathcal{M}_k \times \emptyset \dots \times \emptyset}_{k - position} \times \{f_k\} \ , \quad 1 \leq k \leq m$$

Step 2: The topological covers $\mathcal{W}$, $\mathcal{W}_{\mathbb{B}}$ & $\mathcal{Q}$ of $\mathcal{E}$.

Paracompactness & product structure of $\mathcal{E}$ imply that there exist natural covers constituted of product charts of $\mathcal{A}_{\mathcal{M}}$ & $\mathcal{A}_{\mathcal{F}}$:

If $p \in \mathcal{E}$, choose a $(U; x^1, \dots, x^n)$ at $\pi(p)$ and a (V, y) at $pr_{\mathcal{F}}(p)$. Thus, the product

$$(W ; \{x^A\}_{1 \leq A \leq n+1}) \equiv \{(U \times V ; \{x^a\}_{1 \leq a \leq n} \times y) \in \mathcal{A}_{\mathcal{E}}$$

is a patch at p .

At each $p \in \mathcal{E}$ choose $N + 1$ distinct patches $(W_{i_a}, \chi_{(i_a)})$, $0 \leq a \leq N$. The number $N \in \mathbb{N}$ will be specified later as will. Such arbitrarily but finite many patches with different coordinate functions can be easily illustrated to exist on $\mathcal{M}$

& $\mathcal{F}$: The U -patches of $\mathcal{M}$ can be chosen distinct with "geodesical-coordinates" arguments (diagonalizing $g_{\mathcal{M}}|_U$ in different coordinates, if we choose different initial curves with respect to which we construct the geodesical orbit of the patch). The V -patches of the 1-manifold $\mathcal{F}$ can be chosen using the induced structure of an analytic embedding of $\mathcal{F}$ into some $\mathcal{G}$ of higher dimension restricted on the curve $\mathcal{F}$.

Consider the restriction of each W_{i_a} on their common intersection

$$W_{i_a} \equiv \bigcap_{b=0}^N W_{i_b}, \quad a = 0, 1, \dots, N$$

Thus, we obtain $(N + 1)$ -patches at p covering the same domain on $\mathcal{E}$, with different coordinate functions. Put N of them together to make an N -element class of patches at each $p \in \mathcal{E}$:

$$[W_i] := \{((W_{i_a}, \chi_{(i_a)})) \mid a = 1, \dots, N\}$$

Use now a Euclidean transfer to identify the distinct points $\chi_{(i_1)}(p), \dots, \chi_{(i_N)}(p)$ as the origin $0 \in \mathbb{R}^{n+1}$. Consider the intersection $\bigcap_{a=1}^N \chi_{(i_a)}(W_{i_a})$, which is an open set in $\mathbb{R}^{n+1}$. Within this set lives an open $(n + 1)$ -ball $\mathbb{B}(0, R_i)$ of maximum radius $R_i > 0$. Choose an $r_i < R_i$. Then

$$\mathbb{B}(0, r_i) \subseteq \bigcap_{a=1}^N \chi_{(i_a)}(W_{i_a})$$

Consider now the inversion of $\mathbb{B}(0, r_i)$ via 1 of the coordinates, say the 1st, $\chi_{(i_1)}|_{\mathbb{B}(0, r_i)}$. This implies that within each W_{i_a} lives an analytically diffeomorphic copy of the ball $\mathbb{B}(0, r_i)$:

$$\chi_{(i_1)}^{-1}(\mathbb{B}(0, r_i)) \subseteq W_{i_a} = \bigcap_{b=1}^N \chi_{(i_b)}(W_{i_b}), \quad a = 1, \dots, N$$

We may also define a class of N -elements of inversed balls at each $p \in \mathcal{E}$, distinguished by the different induced coordinates on each of them:

$$[\mathbb{B}_i] := \{(\chi_{(i_1)}^{-1}(\mathbb{B}(0, r_i)), \chi_{(i_a)}|_{\chi_{(i_1)}^{-1}(\mathbb{B}(0, r_i))}) \mid a = 1, \dots, N\}$$

Take now the remaining W_{i_0} patch at p not included in $[W_i]$ and restrict it on $\chi_{i_1}^{-1}(\mathbb{B}(0, r_i))$:

$$(W_{i_0} \equiv W_{i_0}|_{\chi_{i_1}^{-1}(\mathbb{B}(0, r_i))}; \chi_{(i_0)}|_{\chi_{i_1}^{-1}(\mathbb{B}(0, r_i))}) \in \mathcal{A}_{\mathcal{E}}$$

Rename it to Q_j and the coordinate functions to $y_{(j)}^1, \dots, y_{(j)}^{n+1}$:

$$(Q_j, \{y_{(j)}^A\}_{1 \leq A \leq n+1}) := (W_{i_0}|_{\chi_{i_1}^{-1}(\mathbb{B}(0, r_i))}, \chi_{(i_0)}|_{\chi_{i_1}^{-1}(\mathbb{B}(0, r_i))})$$

We form now 3 different topological covers of $\mathcal{E}$, as follows:

Consider the above $(Q_j, \{y_{(j)}^A\}_{1 \leq A \leq n+1})$ patch at each $p \in \mathcal{E}$ & refine to a locally finite cover:

$$\mathcal{Q} := \{(Q_j, \{y_{(j)}^A\}_{1 \leq A \leq n+1}) \mid j \in I\}$$

Dimension theory (see e.g. [9] for an elementary introduction) implies that the *Lebesgue topological dimension* $td(\mathcal{E}) \in \mathbb{N}$ of the paracompact manifold $\mathcal{E}$ equals its geometrical dimension

$$td(\mathcal{E}) = \dim(\mathcal{E}) = n + 1 \in \mathbb{N}$$

This means that no point of $\mathcal{E}$ lies in more than $(n + 1) + 1 = n + 2$ elements of the refinement $\mathcal{Q}$.

The previous cover implies the existence of 2 more locally finite covers, which we obtain considering the classes $[\mathbb{B}_i]$ and $[W_i]$ at each $p \in \mathcal{E}$, refined by the way prescribed by the construction of $\mathcal{Q}$:

$$\mathcal{W}_{\mathbb{B}} := \{[\mathbb{B}_i] \mid i \in \{1, \dots, N\} \times I\} \equiv$$

$$\equiv \{(\chi_{i_1}^{-1}(\mathbb{B}(0, r_i)), \chi_{(i_a)}|_{\chi_{i_1}^{-1}(\mathbb{B}(0, r_i))}) \mid a = 1, \dots, N \mid i_a \in I\}$$

and

$$\mathcal{W} := \{[W_i] \mid i \in \{1, \dots, N\} \times I\} \equiv \{(W_{i_a}; \chi_{(i_a)}) \mid a = 1, \dots, N \mid i_a \in I\}$$

We note that if some $p \in \mathcal{E}$ lies in 1 chart $(Q_j, \{y_{(j)}^A\}_{1 \leq A \leq n+1}) \in \mathcal{Q}$, then it lies in $N - W_{i_a}$'s and $N - \chi_{(i_a)}^{-1}(\mathbb{B}(0, r_i))$'s of $\mathcal{W}$ & $\mathcal{W}_{\mathbb{B}}$ respectively.

Step 3: The family $\{f_{i_a}\}_{i_a \in \{1, \dots, N\} \times I}$ of smooth "Bell" functions on $\mathcal{E}$.

In the sequel we shall need to use the partition functions' tool to obtain a global metric structure. Deviating from the standard approach, we do not introduce a partition subordinate to a locally finite cover, but instead we present constructively a family of functions with similar properties (except for the normalization $\sum f_i = 1$).

This novelty owes to the fact that we will need to use the explicit form of these functions and their crucial property of being real analytic within the interior of their supports.

The "bell functions" characterization owes to the form of their graph.

Statement 2.2. There exists a family $\{f_{i_a}\}_{i_a \in \{1, \dots, N\} \times I}$ of C^∞ non-negative functions on $\mathcal{E}$, with properties:

$$f_{i_a} \in C^\infty(\mathcal{E} \longrightarrow \mathbb{R} \cap [0, +\infty)) \quad \& \quad \text{supp}(f_{i_a}) \subseteq W_{i_a}, \quad \forall [W_i] \in \mathcal{W}$$

$$\text{and} \quad \forall p \in \mathcal{E}, \quad \sum_{i_a \in \{1, \dots, N\} \times I} f_{i_a}(p) > 0$$

and the f_{i_a} 's are real analytic within the set they are strictly positive

$$\{p \in \mathcal{E} / f_{i_a}(p) > 0\} \equiv \{f_{i_a} > 0\}$$

(which is open & coincides with $\text{int}(\text{supp}(f_{i_a})) \equiv \widehat{\text{supp}(f_{i_a})}^\circ$):

$$f_{i_a}|_{\{f_{i_a} > 0\}} = f_{i_a}|_{\widehat{\text{supp}(f_{i_a})}^\circ} \in C_\alpha^\infty(\mathcal{E} \cap \{f_{i_a} > 0\} \longrightarrow \mathbb{R})$$

Remark 2.3. The normalization condition $\sum_{i_a \in \{1, \dots, N\} \times I} f_{i_a}(p) = 1$ is not necessary to be requested, since we do not employ convex functional combinations in the sequel. We are only interested in the existence of a collection of smooth functions, locally analytic within their supports.

Proof. Consider the C^∞ non-negative function on $\mathbb{R}^{n+1}$:

$$f \in C^\infty(\mathbb{R}^{n+1} \longrightarrow [0, +\infty))$$

$$f(x) := \begin{cases} \exp\left(\frac{1}{\|x\|^2 - r^2}\right), & \text{for } \|x\| < r \\ 0, & \text{for } \|x\| \geq r. \end{cases}$$

($r > 0$) which satisfies:

$$\{f > 0\} \equiv \{x \in \mathbb{R}^{n+1} / f(x) > 0\} = \mathbb{B}(0, r) \subsetneq \text{supp}(f)$$

and

$$\text{supp}(f) = \overline{\mathbb{B}(0, r)}$$

and is real analytic in the interior of its support as composition of analytic functions enjoying the same property:

$$f|_{\{f>0\}} = f|_{\mathbb{B}(0, r)} \in C^\infty_\alpha(\mathbb{B}(0, r) \longrightarrow \mathbb{R})$$

(the *Analytic Continuation Principle* implies that f can never be analytic out of $\mathbb{B}(0, r)$, provided that

$$(D^{a_1+\dots+a_s}f)(x) |_{x \rightarrow \partial(\mathbb{B}(0, r))} = 0, \quad \forall a = (a_1, \dots, a_s) \in \mathbb{N}^s$$

see also [10].)

Recall now the cover $\mathcal{W}$ of W -patches and its refinement $\mathcal{W}_\mathbb{B} \subset \mathcal{W}$ of inversed images of open balls and define a family of smooth real functions on (the analytic manifold) $\mathcal{E}$ as:

$$f_{i_a} \in C^\infty(\mathcal{E} \longrightarrow \mathbb{R}), \quad \forall i_a \in \{1, \dots, N\} \times I :$$

$$p \longmapsto f_{i_a}(p) := \begin{cases} \exp\left(\frac{1}{\|\chi_{i_1}(p)\|^2 - r_i^2}\right), & \forall p \in \chi_{i_1}^{-1}(\mathbb{B}(0, r_i)) \subseteq W_{i_a} \\ 0, & \forall p \in \mathcal{E} \setminus \chi_{i_1}^{-1}(\mathbb{B}(0, r_i)). \end{cases} \quad (2.1)$$

Since $\mathcal{E}$ is real analytic, we have

$$f_{i_a}|_{\{f_{i_a}>0\}} = f_{i_a}|_{\chi_{i_a}^{-1}(\mathbb{B}(0,r_i))} \in C_\alpha^\infty(\chi_{i_a}^{-1}(\mathbb{B}(0,r_i))) \longrightarrow \mathbb{R}$$

and the supports of the f_{i_a} 's (as well as their interior) form a locally finite cover of $\mathcal{E}$:

$$\text{supp}(f_{i_a}) = \chi_{i_1}^{-1}(\overline{\mathbb{B}(0,r_i)}) \subseteq W_{i_a},$$

$$\bigcup_{i_a \in \{1,\dots,N\} \times I} \text{supp}(f_{i_a}) = \bigcup_{i_a \in \{1,\dots,N\} \times I} \{f_{i_a} > 0\} = \mathcal{E} \quad \&$$

$$\sum_{i_a \in \{1,\dots,N\} \times I} f_{i_a} > 0$$

and this ends the proof of the *Statement*. $\square$

Step 4: The global smooth & locally analytic metric structure $g_{\mathcal{E}}$ on $\mathcal{E}$.

We can define a global section $g_{\mathcal{E}} \in \Gamma(T^*\mathcal{E} \otimes T^*\mathcal{E})$ as the product metric on $\mathcal{M}_1 \times \dots \times \mathcal{M}_m \times \mathcal{F}$: let $\psi^{(i_a)} \in C_\alpha^\infty(W_{i_a} \longrightarrow \mathbb{R})$ be any analytic function on W_{i_a} (the $\psi^{(i_a)}$'s will be specified later so that this metric to give an Einstein space geometry, using the local 1.3 *Theorem*). First, we define a metric on $\mathcal{M}$ as the product metric on $\mathcal{M}_1 \times \dots \times \mathcal{M}_m$:

$$g_{\mathcal{M}} := g_{\mathcal{M}_1} \circ (dpr_{\mathcal{M}_1} \times dpr_{\mathcal{M}_1}) + \dots + g_{\mathcal{M}_m} \circ (dpr_{\mathcal{M}_m} \times dpr_{\mathcal{M}_m})$$

where $pr_{\mathcal{M}_k} : \mathcal{E} \longrightarrow \mathcal{M}_k$ is the projection on the k -factor. Now, we may set

$$g_{\mathcal{E}} := g_{\mathcal{M}} \circ (d\pi \times d\pi) + \left(\sum_{\substack{i_a \in I \\ 1 \leq a \leq N}} f_{i_a} \psi^{(i_a)} dx_{(i_a)}^{n+1} \otimes dx_{(i_a)}^{n+1} \right) \circ (dpr_{\mathcal{F}} \times dpr_{\mathcal{F}})$$

Introducing the functions:

$$\check{\psi}^{(i_a)} := f_{i_a} \psi^{(i_a)} \in C^\infty(\mathcal{E} \longrightarrow \mathbb{R})$$

(each f_{i_a} given by (2.1)) the metric on $\mathcal{E}$ can be rewritten as

$$g_{\mathcal{E}} = g_{\mathcal{M}} \circ (d\pi \times d\pi) + \left(\sum_{\substack{i_a \in I \\ 1 \leq a \leq N}} \check{\psi}^{(i_a)} dx_{(i_a)}^{n+1} \otimes dx_{(i_a)}^{n+1} \right) \circ (dpr_{\mathcal{F}} \times dpr_{\mathcal{F}}) \quad (2.2)$$

or

$$g_{\mathcal{E}} = \sum_{1 \leq k \leq m} g_{\mathcal{M}_k} \circ (dpr_{\mathcal{M}_k} \times dpr_{\mathcal{M}_k}) + \left(\sum_{\substack{i_a \in I \\ 1 \leq a \leq N}} \check{\psi}^{(i_a)} dx_{(i_a)}^{n+1} \otimes dx_{(i_a)}^{n+1} \right) \circ (dpr_{\mathcal{F}} \times dpr_{\mathcal{F}})$$

Recall now the *Fundamental Theorem of Riemannian Geometry* and equip $(\mathcal{E}, g_{\mathcal{E}})$ with the respective unique Levi-Civita linear connection which we denote by $\nabla^{\mathcal{E}}$. Thus, the the triple

$$(\mathcal{E}; \nabla^{\mathcal{E}}, g_{\mathcal{E}})$$

is an analytic semi-Riemannian manifold with smooth (& locally analytic) metric and the respective metric connection.

If $(D; z^A) \in \mathcal{A}_{\mathcal{E}}$ is any chart of $\mathcal{E}$ (not necessarily product), the presentation $g_{\mathcal{E}}$ in the local coordinates of this patch is:

$$g_{\mathcal{E}}|_D = \{g_{\mathcal{M}}|_D(\frac{\partial\pi}{\partial z^A}, \frac{\partial\pi}{\partial z^B}) + \left(\sum_{\substack{i_a \in I \\ 1 \leq a \leq N}} \check{\psi}^{(i_a)} \frac{\partial(x_{(i_a)}^{n+1} \circ pr_{\mathcal{F}})}{\partial z^A} \frac{\partial(x_{(i_a)}^{n+1} \circ pr_{\mathcal{F}})}{\partial z^B} \right)|_D\} dz^A \otimes dz^B$$

having used that

$$d\pi(\frac{\partial}{\partial z^A}) = \frac{\partial\pi}{\partial z^A}$$

and

$$dx_{(i_a)}^{n+1}(dpr_{\mathcal{F}}(\frac{\partial}{\partial z^A})) = d(x_{(i_a)}^{n+1} \circ pr_{\mathcal{F}})(\frac{\partial}{\partial z^A}) = \frac{\partial(x_{(i_a)}^{n+1} \circ pr_{\mathcal{F}})}{\partial z^A}$$

Taking $D = Q_j$, the form of the product patch implies the following diagonalized form of $g_{\mathcal{E}}|_{Q_j}$ in the respective coordinates:

$$\begin{aligned} g_{\mathcal{E}}|_{Q_j} &= \{g_{\mathcal{M}}(\frac{\partial\pi}{\partial y_{(j)}^a}, \frac{\partial\pi}{\partial y_{(j)}^b})|_{Q_j}\} dy_{(j)}^a \otimes dy_{(j)}^b + \\ &+ \left\{ \left(\sum_{\substack{i_a \in I \\ 1 \leq a \leq N}} \check{\psi}^{(i_a)} \left(\frac{\partial(x_{(i_a)}^{n+1} \circ pr_{\mathcal{F}})}{\partial y_{(j)}^{n+1}} \right)^2 \right) |_{Q_j} \right\} dy_{(j)}^{n+1} \otimes dy_{(j)}^{n+1}, \quad 1 \leq a, b \leq n \end{aligned}$$

for all Q_j 's in $\mathcal{Q}$. Thus, the components of $g_{\mathcal{E}}|_{Q_j}$ are

$$[g_{\mathcal{E}}^{(j)}]_{ab} := g_{\mathcal{M}}\left(\frac{\partial\pi}{\partial y_{(j)}^a}, \frac{\partial\pi}{\partial y_{(j)}^b}\right)|_{Q_j}, \quad 1 \leq a, b \leq n \quad (2.3)$$

and

$$[g_{\mathcal{E}}^{(j)}]_{(n+1)(n+1)} := \left\{ \sum_{\substack{i_a \in I \\ 1 \leq a \leq N}} \check{\psi}^{(i_a)} \left(\frac{\partial(x_{(i_a)}^{n+1} \circ pr_{\mathcal{F}})}{\partial y_{(j)}^{n+1}} \right)^2 \right\}|_{Q_j} \quad (2.4)$$

and they are all analytic functions, under the assumption for the $\psi^{(i_a)}$'s:

$$[g_{\mathcal{E}}^{(j)}]_{(n+1)(n+1)} \quad \& \quad [g_{\mathcal{E}}^{(j)}]_{ab} \in C_{\alpha}^{\infty}(Q_j \longrightarrow \mathbb{R})$$

Step 5: The requested Curvature condition on $\mathcal{E}$ expressed in terms of local coordinates of the cover $\mathcal{Q}$ using the local results.

The condition $\mathcal{E}$ to be an $n + 1$ -dimensional Einstein manifold, with respect to the semi-Riemannian structure $(\nabla^{\mathcal{E}}, g_{\mathcal{E}})$ (Def. 1.2)

$$Ric^{\nabla^{\mathcal{E}}}(g_{\mathcal{E}}) = \frac{2\Lambda}{n-1} g_{\mathcal{E}} \quad (2.5)$$

is equivalent to the condition this equation to hold within all the local coordinated open submanifolds of a cover of $\mathcal{E}$, say $\mathcal{Q}$:

$$Ric^{\nabla^{\mathcal{E}}|_{Q_j}}(g_{\mathcal{E}}|_{Q_j}) = \frac{2\Lambda}{n-1} (g_{\mathcal{E}}|_{Q_j}), \quad \forall j \in I \quad (2.6)$$

and in the local coordinates $\{y_{(j)}^A\}_{1 \leq A \leq n+1}$ of each Q_j the previous relation reads

$$[R_{\mathcal{E}}^{(j)}]_{AB} dy_{(j)}^A \otimes dy_{(j)}^B = \frac{2\Lambda}{n-1} [g_{\mathcal{E}}^{(j)}]_{AB} dy_{(j)}^A \otimes dy_{(j)}^B, \quad \forall j \in I$$

where $[g_{\mathcal{E}}^{(j)}]_{AB}$ are given by (2.3) & (2.4) and $[R_{\mathcal{E}}^{(j)}]_{AB}$ are the components of the Ricci tensor of curvature, calculated by (due to the vanishing *Torsion* of $\nabla^{\mathcal{E}}$):

$$[R_{\mathcal{E}}^{(j)}]_{AB} = \partial_M \Gamma_{(j)AB}^M + \Gamma_{(j)AB}^M \Gamma_{(j)MN}^N - \partial_B \Gamma_{(j)AM}^M - \Gamma_{(j)AN}^M \Gamma_{(j)MB}^N$$

and

$$Ric^{\nabla^\mathcal{E}|_{Q_j}}(g_\mathcal{E}|_{Q_j}) = Ric^{\nabla^\mathcal{E}|_{Q_j}}\left(\frac{\partial}{\partial y_{(j)}^A}, \frac{\partial}{\partial y_{(j)}^B}\right) dy_{(j)}^A \otimes dy_{(j)}^B \equiv [R_\mathcal{E}^{(j)}]_{AB} dy_{(j)}^A \otimes dy_{(j)}^B$$

where the Christoffel symbols are given by

$$\Gamma_{(j)AB}^C = \frac{1}{2} [g_\mathcal{E}^{(j)}]^{CM} \left(-\partial_M([g_\mathcal{E}^{(j)}]_{AB}) + \partial_A([g_\mathcal{E}^{(j)}]_{BM}) + \partial_B([g_\mathcal{E}^{(j)}]_{MA}) \right)$$

Corollary 1.4 implies that, since the functions $[g_\mathcal{E}^{(j)}]_{ab}$ of (2.3) are analytic, there do exist analytic functions $[g_\mathcal{E}^{(j)}]_{(n+1)(n+1)}$, $\forall j \in I$ (at 1 point at least, say $0 \equiv y_{(j)}^A(p) \in \mathbb{R}^{n+1}$) such that equation (2.6) to hold on every Q_j .

Remark 2.4. *The objective pursued to the rest of the proof is to show that $g_\mathcal{E}$ can indeed be fixed through the undetermined $\psi^{(i_a)}$'s living in the W_{i_a} 's of the underlying cover $\mathcal{W}$, so that the local embeddings on every Q_j of $\mathcal{Q}$ to hold, and therefore the whole $\mathcal{E}$ to be an Einstein manifold.*

Step 6: Fixing the Einstein space metric $g_\mathcal{E}$ through the unspecified functions $\psi^{(i_a)} \in C_\alpha^\infty(W_{i_a} \longrightarrow \mathbb{R})$.

The topological dimensionality arguments of *Step 2* imply that no point of $\mathcal{E}$ lies in more than $n + 2$ charts $(Q_j, \{y_{(j)}^A\}_{1 \leq A \leq n+1})$ of $\mathcal{Q}$.

On every Q_j relation (2.4) gives an equation that will be used to determine the $\psi^{(i_a)}$'s in terms of the $[g_\mathcal{E}^{(j)}]_{(n+1)(n+1)}$'s.

$$\underbrace{[g_\mathcal{E}^{(j)}]_{(n+1)(n+1)}}_{\Phi_{(j)}} = \underbrace{\left\{ \sum_{\substack{i_a \in I \\ 1 \leq a \leq N}} f_{i_a} \left(\frac{\partial(x_{(i_a)}^{n+1} \circ pr_\mathcal{F})}{\partial y_{(j)}^{n+1}} \right)^2 \right\}|_{Q_j}}_{\Theta_{(j)}^{(i_a)}} \psi^{(i_a)}$$

This is a linear functional equation with analytic coefficients:

$$\Phi_{(j)} = \sum_{i_a \in I} \{ \Theta_{(j)}^{(i_1)} \psi^{(i_1)} + \dots + \Theta_{(j)}^{(i_a)} \psi^{(i_a)} + \dots + \Theta_{(j)}^{(i_N)} \psi^{(i_N)} \} \quad (2.7)$$

and the $\psi^{(i_a)}$ -variables are all linearly independent, since the coordinates of the patches are chosen distinct, although in the coefficients there exist the same

f_{i_a} 's. Notwithstanding, these functions are distinguished by the different local presentations on the patches.

Consider the domain of $\mathcal{E}$ that appear the maximum $n + 2$ finite-many overlaps of the Q_j 's of the $\mathcal{Q}$ -cover:

$$\bigcap_{r=1}^{n+2} Q_{j_r} = Q_{j_1} \cap \dots \cap Q_{j_{n+2}} \subseteq \mathcal{E}$$

If we wish eq. (2.6) to hold on this intersection, we must request the equations (2.7) to hold on all the Q_j 's simultaneously. This gives rise to a system of linear (in the $\psi^{(i_a)}$'s) equations

$$(\Sigma 1) \quad \begin{cases} \Phi_{(j_1)} = \sum_{i_a=1}^{n+2} \{\Theta_{(j_1)}^{(i_1)} \psi^{(i_1)} + \dots + \Theta_{(j_1)}^{(i_a)} \psi^{(i_a)} + \dots + \Theta_{(j_1)}^{(i_N)} \psi^{(i_N)}\} \\ \vdots \\ \Phi_{(j_{n+2})} = \sum_{i_a=1}^{n+2} \{\Theta_{(j_{n+2})}^{(i_1)} \psi^{(i_1)} + \dots + \Theta_{(j_{n+2})}^{(i_a)} \psi^{(i_a)} + \dots + \Theta_{(j_{n+2})}^{(i_N)} \psi^{(i_N)}\} \end{cases}$$

which admits infinite N -tuples of solutions $\psi^{(i_a)}$ on each $Q_j \cap \bigcap_{r'=1}^{n+2} Q_{j_{r'}}$, if we set $N \geq 2$.

We wish to extend the $\psi^{(i_a)}$ -solutions outside of $\bigcap_{r=1}^{n+2} Q_{j_r}$ on $\bigcup_{r=1}^{n+2} Q_{j_r}$ and finally on the whole $\mathcal{E}$. We follow a stepwise procedure, to illustrate the method.

First, we extend the $\psi^{(i_a)}$'s on the sets

$$Q_{j_2} \cap \dots \cap Q_{j_{n+2}}, \quad \dots, \quad Q_{j_1} \cap \dots \cap \widehat{Q_{j_s}} \cap \dots \cap Q_{j_{n+2}}, \quad \dots, \\ Q_{j_1} \cap \dots \cap Q_{j_{n+1}}$$

where the $(\widehat{})$ -notation indicates that the Q_{j_s} is omitted. These are

$$\binom{n+2}{n+1} = \frac{(n+1)! (n+2)}{(n+1)!} = n+2$$

many sets. If we wish the $\psi^{(i_a)}$'s to be definable on these, we must additionally to $(\Sigma 1)$ request the following equations to hold (setting any sufficient value $N \in \mathbb{N}$ so that to have more $\psi^{(i_a)}$'s than equations)

$$(\Sigma 2) \left\{ \begin{array}{l} \Phi_{(j_1)} = \sum_{\substack{i_a=1 \\ [\mathbb{B}_i] \not\subseteq Q_{j_s}}}^{n+2} \{ \Theta_{(j_1)}^{(i_1)} \psi^{(i_1)} + \dots + \Theta_{(j_1)}^{(i_a)} \psi^{(i_a)} + \dots + \Theta_{(j_1)}^{(i_N)} \psi^{(i_N)} \} \\ \vdots \\ \Phi_{(j_{s-1})} = \sum_{\substack{i_a=1 \\ [\mathbb{B}_i] \not\subseteq Q_{j_s}}}^{n+2} \{ \Theta_{(j_{s-1})}^{(i_1)} \psi^{(i_1)} + \dots + \Theta_{(j_{s-1})}^{(i_a)} \psi^{(i_a)} + \dots + \Theta_{(j_{s-1})}^{(i_N)} \psi^{(i_N)} \} \\ \Phi_{(j_{s+1})} = \sum_{\substack{i_a=1 \\ [\mathbb{B}_i] \not\subseteq Q_{j_s}}}^{n+2} \{ \Theta_{(j_{s+1})}^{(i_1)} \psi^{(i_1)} + \dots + \Theta_{(j_{s+1})}^{(i_a)} \psi^{(i_a)} + \dots + \Theta_{(j_{s+1})}^{(i_N)} \psi^{(i_N)} \} \\ \vdots \\ \Phi_{(j_{n+2})} = \sum_{\substack{i_a=1 \\ [\mathbb{B}_i] \not\subseteq Q_{j_s}}}^{n+2} \{ \Theta_{(j_{n+2})}^{(i_1)} \psi^{(i_1)} + \dots + \Theta_{(j_{n+2})}^{(i_a)} \psi^{(i_a)} + \dots + \Theta_{(j_{n+2})}^{(i_N)} \psi^{(i_N)} \} \end{array} \right.$$

The extension of solutions $\psi^{(i_a)}$ from $\bigcap_{r=1}^{n+2} Q_{j_r}$ to $Q_{j_1} \cap \dots \cap \widehat{Q_{j_s}} \cap \dots \cap Q_{j_{n+2}}$ ($1 \leq s \leq n+2$) has the following meaning:

Requesting the system $(\Sigma 1)$ to hold on $\bigcap_{r=1}^{n+2} Q_{j_r}$, nothing guarantees that there exist solutions $\psi^{(i_a)}$ given by $(\Sigma 2)$ on the $Q_{j_1} \cap \dots \cap \widehat{Q_{j_s}} \cap \dots \cap Q_{j_{n+2}}$'s, which, if restricted, coincide with that of $\bigcap_{r=1}^{n+2} Q_{j_r}$. Consequently, requesting both $(\Sigma 1)$ & $(\Sigma 2)$ to hold on $\bigcap_{r=1}^{n+2} Q_{j_r}$ and merely $(\Sigma 2)$ on $Q_{j_1} \cap \dots \cap \widehat{Q_{j_s}} \cap \dots \cap Q_{j_{n+2}}$, we obtain the requested compatibility of solutions.

Continuing, we can extend the $\psi^{(i_a)}$'s on the $\binom{n+1}{n} = \frac{1}{2}(n+1)(n+2)$ sets

$$Q_{j_3} \cap \dots \cap Q_{j_{n+2}} \ , \ \dots \ ,$$

$$Q_{j_1} \cap \dots \cap \widehat{Q_{j_s}} \cap \dots \cap \widehat{Q_{j_k}} \cap \dots \cap Q_{j_{n+2}} \ , \ \dots$$

$$\dots \ , \ Q_{j_1} \cap \dots \cap Q_{j_n}$$

where $1 \leq s < k \leq n+2$, requesting the additional equations

$$(\Sigma 3) \quad \left\{ \begin{array}{l} \Phi_{(j_r)} = \sum_{\substack{i_a=1 \\ [\mathbb{B}_i] \not\subseteq Q_{j_s} \cup Q_{j_k}}}^{n+2} \{ \Theta_{(j_r)}^{(i_1)} \psi^{(i_1)} + \dots + \Theta_{(j_r)}^{(i_a)} \psi^{(i_a)} + \dots + \Theta_{(j_r)}^{(i_N)} \psi^{(i_N)} \} \\ 1 \leq r \leq n+2, \quad r \neq s, k \end{array} \right.$$

Continuing inductively, we can determine the $\psi^{(i_a)}$'s on $Q_{j_1}, \dots, Q_{j_{n+2}}$, and thus on the whole $Q_{j_1} \cup \dots \cup Q_{j_{n+2}}$.

Hence, the final set of equations we request to hold so that to extend the solutions as previously described is

$$(\Sigma 1) \quad \left\{ \begin{array}{l} \Phi_{(j_r)} = \sum_{i_a=1}^{n+2} \{ \Theta_{(j_r)}^{(i_1)} \psi^{(i_1)} + \dots + \Theta_{(j_r)}^{(i_a)} \psi^{(i_a)} + \dots + \Theta_{(j_r)}^{(i_N)} \psi^{(i_N)} \} \\ 1 \leq r \leq n+2 \end{array} \right.$$

on $Q_{j_1} \cap \dots \cap Q_{j_{n+2}}$

$$(\Sigma 2) \quad \left\{ \begin{array}{l} \Phi_{(j_r)} = \sum_{\substack{i_a=1 \\ [\mathbb{B}_i] \not\subseteq Q_{j_s}}}^{n+2} \{ \Theta_{(j_r)}^{(i_1)} \psi^{(i_1)} + \dots + \Theta_{(j_r)}^{(i_a)} \psi^{(i_a)} + \dots + \Theta_{(j_r)}^{(i_N)} \psi^{(i_N)} \} \\ 1 \leq r \neq s \leq n+2 \end{array} \right.$$

on $Q_{j_1} \cap \dots \cap \widehat{Q_{j_s}} \cap \dots \cap Q_{j_{n+2}}$

$$(\Sigma 3) \quad \left\{ \begin{array}{l} \Phi_{(j_r)} = \sum_{\substack{i_a=1 \\ [\mathbb{B}_i] \not\subseteq Q_{j_s} \cup Q_{j_k}}}^{n+2} \{ \Theta_{(j_r)}^{(i_1)} \psi^{(i_1)} + \dots + \Theta_{(j_r)}^{(i_a)} \psi^{(i_a)} + \dots + \Theta_{(j_r)}^{(i_N)} \psi^{(i_N)} \} \\ 1 \leq r \leq n+2, \quad r \neq s, k \end{array} \right.$$

on $Q_{j_1} \cap \dots \cap \widehat{Q_{j_s}} \cap \dots \cap \widehat{Q_{j_k}} \cap \dots \cap Q_{j_{n+2}}$

$\vdots$
 $\vdots$

$$(\Sigma(n+2)) \quad \left\{ \begin{array}{l} \Phi_{(j_r)} = \sum_{i_a=1}^{n+2} \{ \Theta_{(j_r)}^{(i_1)} \psi^{(i_1)} + \dots + \Theta_{(j_r)}^{(i_a)} \psi^{(i_a)} + \dots + \Theta_{(j_r)}^{(i_N)} \psi^{(i_N)} \} \\ [\mathbb{B}_i] \not\subseteq Q_{j_1} \cap \dots \cap \widehat{Q_{j_r}} \cap \dots \cap Q_{j_{n+2}} \\ r = 1, \dots, n+2 \end{array} \right.$$

on $Q_{j_1}, \dots, Q_{j_{n+2}}$.

The total number of all the equations $(\Sigma 1), (\Sigma 2), \dots, (\Sigma(n+2))$ is $M = \binom{n+2}{n+2} + \binom{n+2}{n+1} + \dots + \binom{n+2}{1}$, that is

$$M = 2n + 5 + \sum_{t=2}^n \frac{(n+3-t) \cdots (n+2)}{t!}$$

or

$$M = 2\left(\sum_{k=1}^m n_k\right) + 5 + \sum_{t=2}^{\sum_{k=1}^m n_k} \frac{\left(\sum_{k=1}^m n_k + 3 - t\right) \cdots \left(\sum_{k=1}^m n_k + 2\right)}{t!}$$

Thus, setting

$$N \equiv M + 1$$

we obtain more $\psi^{(i_a)}$'s than equations, thus, there do exist solutions as desired so that the curvature condition given by *eq.* (2.6) to hold on the union $Q_{j_1} \cup \dots \cup Q_{j_{n+2}}$.

We claim that in fact we have specified the metric $g_{\mathcal{E}}$ through the $\psi^{(i_a)}$'s on the whole $\mathcal{E}$. If $\bigcup Q_j, \bigcup Q_{j'}$ are 2 unions of Q_j 's overlapping on at least 1 Q_{j_0} , then the metrics $g_{\mathcal{E}}|_{\bigcup Q_j}, g_{\mathcal{E}}|_{\bigcup Q_{j'}}$ coincide on $\bigcup Q_{j'} \cap \bigcup Q_j \subseteq Q_{j_0}$, provided that we have requested the equation

$$\Phi_{(j_0)} = \sum_{\substack{i_a=1 \\ [\mathbb{B}_i] \subseteq Q_{j_0}}}^{n+2} \{\Theta_{(j_0)}^{(i_1)} \psi^{(i_1)} + \dots + \Theta_{(j_0)}^{(i_a)} \psi^{(i_a)} + \dots + \Theta_{(j_0)}^{(i_N)} \psi^{(i_N)}\}$$

to hold on both systems solved on $\bigcup Q_j$ and $\bigcup Q_{j'}$.

Step 7: The isometry & C_α^∞ embedding conditions.

Having shown that $g_{\mathcal{E}}$ given by (2.2) does exist so that relation (2.5) to hold on $\mathcal{E}$, the isometry condition holds trivially by restriction of the product metric on the submanifolds $\mathcal{M}_k$:

$$g_{\mathcal{E}}|_{\mathcal{M}_k} = g_{\mathcal{M}_k} \circ (dpr_{\mathcal{M}_k} \times dpr_{\mathcal{M}_k}) \equiv g_{\mathcal{M}_k} \in \Gamma(T^*\mathcal{M}_k \otimes T^*\mathcal{M}_k)$$

if any $\xi, \eta \in T\mathcal{M}_k$ are applied. The Riemannian connection $\nabla^{\mathcal{E}}$ of $\mathcal{E}$ restricted on (the open in the induced subspace topology) $\mathcal{M}_k$'s reduce to the same $\nabla^{\mathcal{M}_k}$

on $\mathcal{M}_k$

$$\nabla^{\mathcal{E}}|_{\mathcal{M}_k} = \nabla^{\mathcal{M}_k}$$

which is the Riemannian connection $\nabla^{\mathcal{M}_k}$ of $\mathcal{M}_k$ with respect to $g_{\mathcal{M}}$. The trivial C^∞_α -analyticity of the inclusion maps $i_{\mathcal{M}_k} : \mathcal{M}_k \hookrightarrow \mathcal{E}$ clearly completes the proof. $\square$

3. Homotopy of Analytic Einstein Global Embedding Manifolds.

In this section we study Homotopically the possible topologies that the embedding space $\mathcal{E}$ can have, in complete analogy with the respective of [26].

The key-property is not the geometry of $\mathcal{E}$ presented in *Th.* 2.1, but the product structure of $\mathcal{E}$.

Proposition 3.1. (*Homotopy of the embedding Einstein manifold*)

The Homotopy of the embedding Einstein space $(\mathcal{E}; \nabla^{\mathcal{E}}, g_{\mathcal{E}})$ of $\mathcal{M}_1, \dots, \mathcal{M}_m$ splits in every dimension as

$$\pi_k(\mathcal{E}) \cong \pi_k(\mathcal{M}_1) \bigoplus \dots \bigoplus \pi_k(\mathcal{M}_m) \bigoplus \pi_k(\mathcal{F}) \quad (3.8)$$

for all $k \in \mathbb{N}$ and any analytic manifold $\mathcal{F}$, $\dim(\mathcal{F}) \geq 1$.

Proof. The proof is a triviality. We simply recall that the bulk $\mathcal{E}$ was defined as the product

$$\mathcal{E} = \mathcal{M}_1 \times \dots \times \mathcal{M}_m \times \mathcal{F}$$

and classical Homotopy theory (e.g. [1], [6], [13]) implies the requested splitting relation for the Homotopy of $\mathcal{E}$. $\square$

Corollary 3.2. *If we choose in the case of the embedding codimension 1, the fibre $\mathcal{F}$ to be homotopically equivalent to $\mathbb{R}$ (or the relatively compact $(0, 1)$) or the circle S^1 :*

$$\mathcal{F} \simeq \mathbb{R} \quad \text{or} \quad \mathcal{F} \simeq S^1$$

we obtain the following respective Homotopies for $\mathcal{E}$

$$\pi_k(\mathcal{E}) \cong \pi_k(\mathcal{M}), \quad \forall k \in \mathbb{N}$$

or

$$\pi_k(\mathcal{E}) \cong \begin{cases} \pi_1(\mathcal{M}_1) \oplus \cdots \oplus \pi_1(\mathcal{M}_m) \oplus \mathbb{Z}, & \text{for } k = 1 \\ \pi_k(\mathcal{M}_1) \oplus \cdots \oplus \pi_k(\mathcal{M}_m), & \forall k \in \mathbb{N} \setminus \{1\}. \end{cases}$$

Remark 3.3. In the proof of 2.1, $\mathcal{E}$ was defined as a product manifold for technical reasons, providing a bulk containing the given $\mathcal{M}$.

Notwithstanding, the direct sum decomposition of its Homotopy implies the *existence* of "simple" topologies for $\mathcal{E}$, with respect to those that the $\mathcal{M}_k$'s may admit.

Acknowledgement. *The Author is profoundly indebted to professor Lorenz Magaard at University of Honolulu, Hawaii for his interest in the present work, P. Krikelis & A. Kartsaklis at University of Athens for helpful comments & technical support and Ed. Anderson for discussions on the physical aspects of the embedding results.*

References

- [1] Glen E. Bredon, *Topology and Geometry*, Springer, 1997
- [2] J. E. Campbell, *A Course of Differential Geometry*, Oxford, Clarendon Press, 1923
- [3] Do Carmo, M.P., *Riemannian Geometry*, Birkhäuser, 1992
- [4] L. Pf. Eisenhart, *Riemannian Geometry*, Princeton University Press, 1964
- [5] O. Gron and S. Hervik, *Einstein's General Theory of Relativity* (e-book), <http://www.fys.uio.no/~sigbjorh/gravity.pdf>, 2004
- [6] A. Hatcher, *Algebraic Topology*, <http://www.math.cornell.edu/~hatcher>, (e-book), 2000

- [7] S. W. Hawking & G. F. R. Ellis *The Large scale structure of spacetime*, Cambridge University Press, 1973
- [8] L. Magaard, *Zur Einbettung Riemannscher Raume in Einstein - Raume und Konfor - Euclidische Raume*, PhD thesis, (Kiel), 1963
- [9] J. R. Munkres, *Topology, a first course*, Prentice-Hall, Inc., Englewood Cliffs, New Jersey
- [10] R. Narasimhan, *Analysis on Real and Complex Manifolds*, Masson and Cie, Paris, North Holland, 1968
- [11] R. Penrose, *Techniques of Differential Topology in Relativity*, SIAM Publ. Philadelphia, 1972
- [12] R. K. Sachs and H. Wu, *General Relativity for Mathematicians*, Springer-Verlang, 1977
- [13] N. Steenrod, *The topology of fibre Bundles*, Princeton University Press, 1972
- [14] H. Stefani, *General Relativity*, 2nd Edition, Cambridge University Press, 1990
- [15] P. S. Wesson, *Space-Time-Matter, Modern Kaluza-Klein Theory*, World Scientific, 2000
(Articles)
- [16] E. Anderson, J. E. Lidsey, *Embeddings in NonVacuum Spacetimes*, Class.Quant.Grav. 18 (2001) 4831-4844, ArXiv: gr-qc/0106090
- [17] F. Dahia, C. Romero, *The embedding of spacetime in 5 dimensions: An extension of Campbell-Magaard theorem*, J.Math.Phys. 43 (2002) 5804-5814, ArXiv: gr-qc/0109076
- [18] F. Dahia, C. Romero, *On the embedding of branes in five-dimensional spaces*, Class.Quant.Grav. (2004) 21, No 4, 927-933, ArXiv: gr-qc/0308056

- [19] A. Davidson, $\Lambda = 0$ cosmology of a brane-like universe, Class. Quantum Grav. 16 (1999) 653–659
- [20] C. Deffayet, *Cosmology of Brane-Induced Gravity Models*, International Journal of Theoretical Physics, Vol. 41, No. 11, November 2002 <http://www.kluweronline.com/article>
- [21] M. Gogberashvili, *Brane-Universe in Six Dimensions with Two Times*, Phys.Lett. B484 (2000) 124-128, ArXiv: hep-ph/0001109
- [22] M. Gogberashvili, *Embedding of the Brane into Six Dimensions*, J.Math.Phys. 43 (2002) 4886–4888, ArXiv: gr-qc/0202061
- [23] M. Gogberashvili, *Brane Gravity from Bulk Vector Field*, Phys.Lett. B553 (2003) 284-288, ArXiv: hep-th/0203090
- [24] H. Grauert, *On Levi's problem and the imbedding of real analytic manifolds*, Annals of Mathematics, 68 (47-82)
- [25] E. Gravanis, S. Willison, *Israel conditions for the Gauss-Bonnet theory and the Friedmann equation on the brane universe*, ArXiv: hep-th/0209076
- [26] N. I. Katzourakis "The Global Embedding Problem of Semi- Riemannian into Einstein Manifolds", ArXiv: math-ph/0407067
- [27] N. I. Katzourakis "On the Global Embedding of Spacetime into Singular $\mathcal{E}^\Sigma$ Einstein Manifolds: Wormholes", ArXiv: math.DG/0503129
- [28] D. Langlois, *Brane cosmology: an introduction*, Prog.Theor.Phys.Suppl. 148 (2003) 181-212, ArXiv: hep-th/0209261
- [29] D. Langlois, *Gravitation and Cosmology in Brane-Worlds*, (Invited plenary talk at the GR17 conference, Dublin, Ireland, July 2004), ArXiv: gr-qc/0410129
- [30] J. E. Lidsey, C. Romero, R. Tavakol, S. Rippl, *On Applications of Campbell's Embedding Theorem*, Class.Quant.Grav. 14 (1997) 865-879, ArXiv: gr-qc/9907040

- [31] J. E. Lidsey, *The Embedding of Superstring Backgrounds in Einstein Gravity*, Phys.Lett. B417 (1998) 33, ArXiv: hep-th/9907095
- [32] R. Maartens, *Geometry and Dynamics of the Brane-World*, Reference Frames and Gravitomagnetism, ed. J Pascual-Sanchez et al. (World Sci., 2001), p. 93-119, ArXiv: gr-qc/0101059
- [33] R. Maartens, *Brane-World Gravity*, Living Rev. Relativity, 7, (2004), 7, <http://www.livingreviews.org/lrr-2004-7>
- [34] M. Pavisic, *Resource Letter on Geometrical Results for Embeddings and Branes*, arXiv: gr-qc/0010045
- [35] A. Perez-Lorenzana, *An Introduction to the Brane World*, ArXiv: hep-ph/0406279
- [36] F. Quevedo, *Lectures on String/Brane Cosmology*, arXiv: hep-th/0210292
- [37] L. Randall, R. Sundrum, *A Large Mass Hierarchy from a Small Extra Dimension*, Phys.Rev.Lett. 83 (1999) 3370-3373, ArXiv: hep-ph/9905221
- [38] L. Randall, R. Sundrum, *An Alternative to Compactification*, Phys.Rev.Lett. 83 (1999) 4690-4693, ArXiv: hep-th/9906064
- [39] S. S. Seahra, *Physics in Higher-Dimensional Manifolds*, <http://etd.uwaterloo.ca/etd/ssseahra2003.pdf>
- [40] S. S. Seahra, P. S. Wesson, *Application of the Campbell- Magaard theorem to higher-dimensional physics*, Class.Quant.Grav. 20 (2003) 1321-1340, ArXiv: gr-qc/0302015
- [41] T. Shiromizu, K. Koyama, *Low energy effective theory for two branes system - Covariant curvature formulation*, Phys.Rev. D67 (2003) 084022, ArXiv: hep-th/0210066
- [42] D. A. Steer, M. F. Parry, *Brane Cosmology, Varying Speed of Light and Inflation in Models With One or More Extra Dimensions*, Int.J.Th.Phys., Vol. 41, No. 11, Nov. 2002, <http://www.kluweronline.com/article>

- [43] D. Wands, *Brane-world cosmology*,
<http://www.tech.port.ac.uk/staffweb/wandsd/powerpoint/sussex.pdf>
- [44] H. Whitney, *Differentiable Manifolds*, Annals of Mathematics, 37 (645-680),
1936